

## Colonoscopy Preparation using *Suprep* 2 Day PM/AM Spilt Dosing Regimen

### • **SEVEN DAYS BEFORE THE PROCEDURE**

#### **Medication restrictions – Please call the prescribing physician**

1. *Diabetics taking insulin:* will need dose adjustment the day before the exam. Please ask for advice from the physician that prescribes your insulin. For diabetics only taking diabetic pills simply stop the pills the day before the procedure (see below).
2. **DO NOT** take ***Ozempic, Mounjaro, Wegovy, Trulicity, Semaglutide, Saxenda, Vitoza, Rybelus, Adlyxin, Byureon and Byetta.*** 1 week prior to procedure, this slows gastric motility and the effectiveness of the prep and increases risk of aspiration if food and liquids are still present in the stomach.
3. *Blood thinners (antithrombotic agents) including: Coumadin (Warfarin), Plavix, Pradaxa, Xarelto, Eliquis, Brilinta and Effient:* These can increase the risk of bleeding during and after endoscopic procedures to a varying degree. **If you take a blood thinner, review our recommendations below for holding these medications prior to your procedure and confirm with your prescribing provider:**

**Effient - 7 days; Coumadin (Warfarin), Plavix and Brilinta - 5 days; Pradaxa, Eliquis and Xarelto - 2 days;**

**\*\* IF these medications are unable to be stopped, you need to call our office ASAP\*\***

4. Aspirin does **NOT** need to be stopped

#### **Items to Purchase**

1. Fill the prescription for the *Suprep* (bowel prep kit).
2. Fill the prescription for *Ondansetron* (anti-nausea medication).
3. Any over-the-counter hemorrhoid ointment for anal discomfort (if needed).
4. Flavored water (i.e. Propel or Nestle Splash), (optional) recommend berry flavor and must be non-carbonated, will need at least 20 ounces. The *Suprep* is berry flavored so any other flavor of may be less appealing.

#### **Diet restrictions**

No seeds, nuts, popcorn or peels from fruits and vegetables. Examples but not limited to: no peels and seeds from apple, potatoes, cucumbers, zucchini, squash, tomatoes, no berries (blueberries, raspberries, strawberries), no grapes or bananas etc.

### • **ONE DAY BEFORE THE PROCEDURE**

1. **For breakfast AND lunch you may choose one of the following:**  
**2 eggs** (no meat, cheese, veggies or toast, just eggs, salt, pepper, oil or butter is fine to cook with)  
**or 1 yogurt, 8 ounces** (plain, can be flavored but no chunks of fruit or granola), **or**  
**1 Ensure, 8 ounces** (any flavor) **then clear liquids only for the remainder of the day.**
2. **You cannot have milk or orange juice for the entire day.**
3. Drink as much as you can to prevent dehydration. You may have bouillon, broth, black coffee (sugar and artificial sweeteners are okay, no creamer of any kind), tea (honey is okay), juice (apple, grape, cranberry), Gatorade, Powerade, lemonade, soda (regular or diet are okay), water, Popsicles and Jell-O (any flavor or color). Taking red Jell-O and red Gatorade may turn your bowel movements red and should be limited. If you have had dehydration before drink Pedialyte.
4. You may have clear candy (such as Lifesavers, Jolly Ranchers and gummy bears).
5. Do not take oral diabetic pills.

**TURN THE PAGE OVER FOR THE REST OF THE DIRECTIONS**

# TIMING OF PREPARATION

**Please follow these instructions and NOT the *Suprep* box instructions**

## • EVENING BEFORE THE PROCEDURE

This is a 2 step process divided into two 16 ounce sessions. You must drink the entire *Suprep* to ensure your colon will be cleaned out. A poor prep can mean missed lesions (flat polyps) and cancellation and rescheduling of your procedure.

**Starting at 6:00pm** (This **CAN NOT** be done earlier in the day.)

1. Take the *Ondansetron* (anti-nausea) 30 minutes prior to taking each dose of the *Suprep* (bowel prep).
2. Pour **ONE** (1) 6 ounce bottle of *Suprep* into the supplied mixing cup; add cold drinking water or berry flavored water to the 16 ounce fill line on the mixing cup and stir. Using the flavored water can help cover up the taste from the laxative.
3. Drink the entire cup of solution. Drinking with straw helps. Drink gradually to prevent nausea.
4. Drink 2 more 16 ounce glasses of water or other clear liquids slowly over the next 1-2 hours (32 ounces total); you can use the supplied mixing cup, filling to fill line on cup.
5. You can continue the clear liquid diet after you finish the 6:00pm dose of the prep.

## • THE MORNING OF THE PROCEDURE

**Beginning 6 hours prior to **scheduled procedure time**, repeat steps 1-4 as above.** (Yes, this requires you to do an early morning or in the middle of the night based on your procedure time.)

1. **Take nothing by mouth, including water, after completing the last *additional 32 ounces of fluid or at least 4 hours prior to procedure time.***
2. You should take any necessary medications you normally use on the morning of the procedure with **sips** of water, vitamins and supplements should be taken after the procedure.
3. Arrive at your scheduled arrival time.
4. Bring a list of your medications, insurance cards and photo ID. If you are a woman of childbearing age you will be required to give a urine specimen when you arrive.
5. You must have a responsible adult drive you home after your procedure due to the sedation. It is best if your driver waits for you at the endoscopy center. **Taking a Cab or an Uber is NOT an option.** If you do not have a driver, we will not be able to sedate you for your procedure and the procedure may be canceled.
6. You may not drive or work until the next day.

## • IMPORTANT INFORMATION ABOUT SEDATION

Sedatives used for endoscopy include Propofol. These medications provide amnesia and pain relief. You are breathing on your own. You may be groggy after your procedure and you **SHOULD NOT** work or drive the rest of the day, a responsible adult must drive you home after your procedure.

## • QUESTIONS OR PROBLEMS:

1. Daytime phone number is (314) 997-0554.
2. After hours number is (314) 388-6578.
3. Internet website address is [www.gidoctor.net](http://www.gidoctor.net)